
Dr. Amit Sachdeva & Nasim Ahmed

¹Assistant Professor, Department of Community Medicine, Indira Gandhi Medical College, Shimla, Himachal Pradesh, India. Email: dramitsachdeva2410@gmail.com

²Independent Research Scholar, Iarcon international LLP, Guwahati, Assam India. Email: nasim@iarcon.org

Received: Jan. 03, 2025; Revised: Feb. 09, 2025; Accepted: Mar. 16, 2025 ; Published: June. 28, 2025

Under a Creative Commons license Doi: 10.47310/ml.2025.v02i01.011

From Malnutrition to Obesity: Public Health Issue of this Double Burden in Indian Paediatric Nutrition

Abstract: India is currently grappling with the "double burden" of malnutrition, where undernutrition and obesity coexist as major public health issues among its pediatric population. While undernutrition remains prevalent in rural and impoverished areas, the rapid rise of childhood obesity in urban regions is driven by unhealthy diets and sedentary lifestyles. This paradox requires a comprehensive public health approach involving government policies, school-based interventions, community efforts, and public awareness campaigns to ensure balanced nutrition and active lifestyles for children.

Keywords: malnutrition, obesity, pediatric health, nutrition, public health

INTRODUCTION

India is facing a paradoxical situation in the realm of paediatric nutrition: while significant strides have been made in combating undernutrition, the country is now witnessing the simultaneous rise of childhood obesity. This coexistence of undernutrition and overnutrition, often referred to as the "double burden" of malnutrition, represents a growing public health challenge. On one hand, a considerable proportion of Indian children suffer from stunted growth, wasting, and micronutrient deficiencies; on the other hand, an increasing number of children are becoming overweight or obese due to the adoption of unhealthy diets and sedentary lifestyles.[1-3]

This dual challenge is exacerbated by India's socio-economic diversity, where malnutrition persists in poorer, rural regions, and obesity is rising in wealthier, urban areas. This article provides an in-depth review of the double burden of malnutrition and obesity in Indian paediatric populations, examining its causes, consequences, and the necessary public health interventions to address this crisis.

The State of Malnutrition in India [4-10]

Current Trends in Undernutrition

Undernutrition continues to be a significant public health concern in India, especially among children under the age of five. Malnutrition manifests in various forms, including stunting (low height for age), wasting (low weight for height), and underweight (low weight for age). These indicators reflect chronic and acute malnutrition that result from inadequate food intake, poor health conditions, and a lack of essential nutrients during critical periods of growth.

Stunting remains one of the most prevalent forms of malnutrition in India, with millions of children unable to reach their full physical and cognitive potential due to long-term nutrient deficiencies. Stunting is particularly common in rural areas, where access to a diverse and nutritious diet is limited. Children who are stunted are not only shorter than their peers but also more vulnerable to illness, poor school performance, and lower productivity in adulthood.

Wasting, another critical concern, is an indicator of acute malnutrition that occurs when children experience rapid weight loss due to insufficient food intake or illness. Wasting often results from

Sachdeva, Amit. "From Malnutrition to Obesity: Public Health Issue of This Double Burden in Indian Paediatric Nutrition." *Medical Letter*, vol. 2, no. 1, 2025, pp. 55–59.

From Malnutrition to Obesity: Public Health Issue of this Double Burden in Indian Paediatric Nutrition

emergencies such as droughts, floods, or conflicts, but it can also be a chronic problem in regions where food insecurity is persistent.

Key Contributing Factors to Malnutrition

1. **Poverty and Food Insecurity:** In rural and impoverished regions, many families lack the resources to provide their children with a balanced and nutritious diet. Limited access to fresh fruits, vegetables, protein sources, and fortified foods leaves children vulnerable to nutrient deficiencies. Additionally, food insecurity, which refers to the lack of consistent access to enough nutritious food, is a significant driver of malnutrition in India. Families may rely on cheap, calorie-dense foods that are low in essential nutrients, exacerbating the issue.
2. **Poor Maternal Health and Care Practices:** Malnutrition often begins in the womb, with many women entering pregnancy in a malnourished state. Poor maternal nutrition, combined with inadequate prenatal care, increases the risk of low birth weight and malnourished infants. After birth, suboptimal breastfeeding practices and the introduction of poor-quality complementary foods further contribute to malnutrition in young children.
3. **Infectious Diseases and Poor Sanitation:** Recurrent infections, such as diarrhoea and respiratory illnesses, are common among malnourished children and further weaken their bodies, preventing the absorption of essential nutrients. Inadequate access to clean water and sanitation also exacerbates the problem, as children who suffer from frequent illnesses struggle to recover their nutritional status.
4. **Lack of Nutrition Education:** Many families, particularly in rural areas, lack awareness of the importance of a balanced diet. Cultural practices, food taboos, and limited knowledge about nutrition contribute to poor feeding practices that deprive children of the variety of nutrients they need for healthy growth and development.

The Rise of Childhood Obesity in India [11-14]

Current Trends in Overnutrition

While malnutrition in the form of undernutrition has long been a public health priority in India, recent years have seen a disturbing rise in childhood obesity, particularly in urban areas. As India undergoes rapid urbanization and economic development, the dietary patterns of its population have shifted dramatically. Children in cities are increasingly exposed to energy-dense, nutrient-poor foods that are high in sugar, fat, and salt. The increased consumption of junk food, sugary beverages, and processed snacks is contributing to rising obesity rates among Indian children.

This shift towards a more "Westernized" diet is compounded by a significant decline in physical activity. Children today spend more time in sedentary activities such as watching television, playing video games, and using smartphones. Schools, particularly in urban areas, often lack adequate spaces for physical education, and parents may prioritize academic success over sports and outdoor play, leading to reduced opportunities for exercise.

Key Contributing Factors to Childhood Obesity

1. **Dietary Shifts:** The proliferation of fast-food outlets, processed snacks, and sugary drinks in urban areas has dramatically altered the dietary habits of Indian children. These foods are often marketed aggressively to children, making them highly appealing. High-calorie, low-nutrient foods are readily available and affordable, leading many children to replace traditional, nutritious meals with unhealthy alternatives. Furthermore, the influence of global food chains and their promotion of "convenience" foods has had a significant impact on the eating patterns of urban families.
2. **Sedentary Lifestyles:** Urbanization has led to a decline in physical activity levels among children. As more children engage in sedentary pastimes, such as screen time on televisions, computers, and mobile devices, the opportunities for physical activity have diminished. In many urban settings, the lack of safe outdoor spaces, such as parks and playgrounds, further restricts children's ability to engage in active play. This sedentary behavior, combined with a high-calorie diet, is a primary contributor to childhood obesity.
3. **Parental Influence and Lifestyle Changes:** In many families, especially in urban areas, both parents may work long hours, leading to a reliance on convenience foods and takeaways. This shift in family dynamics has led to a growing dependence on processed foods and less time for home-cooked, nutritious meals. Moreover, children often mirror the dietary habits of their

From Malnutrition to Obesity: Public Health Issue of this Double Burden in Indian Paediatric Nutrition

parents, and when parents lead sedentary lives and consume unhealthy foods, children are likely to adopt similar behaviors.

4. **Socio-Economic Changes:** As India's middle class grows, the consumption of high-calorie, high-fat foods is often seen as a status symbol. Families may view fast food and packaged snacks as signs of modernity and affluence, further contributing to the rise in childhood obesity. Additionally, with increased disposable income, urban families tend to spend more on convenience foods, which are often less nutritious than traditional home-cooked meals.

The Double Burden of Malnutrition and Obesity [7-12]

India's double burden of malnutrition reflects the country's complex socio-economic landscape, where undernutrition and obesity coexist within the same communities, and even within the same households. Children in rural and impoverished areas continue to suffer from undernutrition, while their counterparts in urban and wealthier families are increasingly at risk of overnutrition. This dual burden poses a unique public health challenge, as the country must simultaneously address issues of food scarcity and food excess.

Nutritional Inequality

One of the key drivers of this double burden is the growing inequality in access to nutritious food. In rural areas, children often lack access to a diverse diet rich in vitamins and minerals. Meanwhile, in urban areas, children may have access to food, but it is often calorie-rich and nutrient-poor, leading to the rise in obesity. This nutritional inequality highlights the need for public health interventions that address both undernutrition and overnutrition, while promoting access to healthy, balanced diets for all children.

Impact on Health Outcomes

The health consequences of the double burden are far-reaching. Children who suffer from undernutrition are at greater risk of infections, cognitive impairments, and developmental delays, while children who are overweight or obese are more likely to develop non-communicable diseases such as diabetes, hypertension, and heart disease at a young age. The coexistence of these two forms of malnutrition increases the overall burden on India's healthcare system, as it must tackle both communicable diseases related to undernutrition and lifestyle-related diseases linked to obesity.

Public Health Interventions and Solutions [3,5,7,12,14]

Addressing the double burden of malnutrition and obesity in India requires a comprehensive, multi-pronged approach that includes government action, community engagement, and family-level interventions.

1. Government Policies and Nutrition Programs

The Indian government has introduced several policies and programs aimed at addressing malnutrition in children, particularly in rural and impoverished areas. These programs focus on improving maternal and child nutrition through targeted interventions such as supplemental feeding, fortification of staple foods, and promoting exclusive breastfeeding. However, to tackle the rising issue of childhood obesity, there is a growing need to expand these initiatives to include education on healthy eating habits, portion control, and the risks of overnutrition.

Government campaigns that promote the consumption of traditional, nutrient-rich foods such as grains, pulses, fruits, and vegetables can help shift dietary patterns away from processed, calorie-dense foods. Additionally, regulatory measures such as imposing taxes on sugary beverages, restricting junk food advertising targeted at children, and mandating clear nutritional labeling on food products could help curb the consumption of unhealthy foods.

2. School-Based Interventions

Schools play a critical role in shaping children's eating habits and lifestyle choices. School meal programs can provide children with nutritious meals that meet their dietary needs, particularly in areas where food insecurity is prevalent. At the same time, schools can promote physical activity through sports programs and physical education classes, encouraging children to lead active lives.

In urban areas, schools can educate children about the importance of a balanced diet and the dangers of excessive junk food consumption. Implementing "no junk food" policies in school canteens, along with nutrition education curricula, can help children make healthier food choices. Schools can also create

From Malnutrition to Obesity: Public Health Issue of this Double Burden in Indian Paediatric Nutrition

environments that encourage physical activity by providing adequate space for sports and recreational activities.

3. Community and Family-Level Interventions

At the community level, local governments and non-governmental organizations can promote access to fresh, nutritious food through initiatives such as community gardens, farmers' markets, and nutrition workshops. These initiatives can help families in both rural and urban areas access affordable, healthy food options while educating them about the benefits of a balanced diet.

Families play a crucial role in preventing both undernutrition and obesity. Parents can model healthy eating habits by preparing balanced, home-cooked meals and limiting their children's intake of processed foods. Encouraging children to participate in meal preparation can also foster an understanding of nutrition and healthy eating. Moreover, families can reduce sedentary behavior by encouraging outdoor play, limiting screen time, and prioritizing physical activities such as walking, cycling, and sports.

4. Public Awareness Campaigns

Raising awareness about the double burden of malnutrition and obesity through mass media campaigns can help shift societal attitudes towards nutrition. Public health campaigns that emphasize the importance of balanced diets, portion control, and physical activity can help reduce the stigma around malnutrition while addressing the growing obesity epidemic. These campaigns should target both urban and rural populations, adapting messages to reflect the unique challenges faced by each group.

CONCLUSION

India's paediatric population is caught in the grips of a double burden of malnutrition, where undernutrition and obesity coexist, each presenting distinct yet interconnected public health challenges. Undernutrition continues to afflict millions of children, particularly in rural and impoverished areas, while the rapid rise of childhood obesity in urban regions threatens the health of future generations.

Addressing this double burden requires a comprehensive approach that tackles the root causes of both undernutrition and obesity. Public health interventions must promote access to healthy, nutritious food while discouraging the consumption of processed, high-calorie foods. Schools, communities, and families must work together to create environments that foster healthy eating habits and active lifestyles for children.

By investing in both preventive and curative strategies, India can protect its children from the devastating health consequences of malnutrition and obesity, ensuring that future generations can thrive and reach their full potential.

REFERENCES

1. Nguyen et al; "The Double Burden of Malnutrition in India: Trends and Inequalities (2006-2016)" 16.2 (2021) Ppe0247856, doi <https://doi.org/10.1371/journal.pone.0247856>
2. Prithishkumar et al; "Double Burden of Malnutrition Among Women of Reproductive Age: Trends and Determinants Over the Last 15 Years in India" 19.6 (2024) Ppe0304776, doi <https://doi.org/10.1371/journal.pone.0304776>
3. Debnath et al; "Double Burden of Malnutrition Among Adolescents in India" 8.2 (2019) Pp155-178
4. World Health Organization et al; "Malnutrition" 1.1 (2021) Pp1-2, Available from: <https://www.who.int/news-room/fact-sheets/detail/malnutrition>
5. Drishti IAS et al; "Combating Malnutrition in India" 1.1 (2023) Pp1-2, Available from: <https://www.drishtiias.com/daily-updates/daily-news-editorials/combating-malnutrition-in-india>
6. Press Information Bureau et al; "Malnutrition-Free India" 1.1 (2021) Pp1-2, Available from: <https://pib.gov.in/PressReleasePage.aspx?PRID=1781673>
7. Feeding India et al; "Understanding the Malnutrition Crisis in India" 1.1 (2023) Pp1-2, Available from: <https://www.feedingindia.org/blog/understanding-the-malnutrition-crisis-in-india>
8. Sahu et al; "Malnutrition Among Under-Five Children in India and Strategies for Control" 6.1 (2015) Pp18-23, doi <https://doi.org/10.4103/0975-7406.154424>

From Malnutrition to Obesity: Public Health Issue of this Double Burden in Indian Paediatric Nutrition

9. *The Times of India et al; "India's Malnutrition Crisis: 17% of Children Underweight, 36% Stunted, 6% Wasted: Reports WCD" 1.1 (2023) Pp1-2, Available from: <https://timesofindia.indiatimes.com/india/indias-malnutrition-crisis-17-of-children-underweight-36-stunted-6-wasted-reports-wcd/articleshow/112045471.cms>*
10. *Next IAS et al; "Malnutrition in India" 1.1 (2022) Pp1-2, Available from: <https://www.nextias.com/ca/current-affairs/28-07-2022/malnutrition-in-india-2>*
11. *Saha et al; "Overweight/Obesity Prevalence Among Under-Five Children and Risk Factors in India: A Cross-Sectional Study Using the National Family Health Survey (2015-2016)" 14.17 (2022) Pp3621, doi <https://doi.org/10.3390/nu14173621>*
12. *Singh et al; "Childhood Obesity in India: A Two-Decade Meta-Analysis of Prevalence and Socioeconomic Correlates" 23.1 (2023) Pp101390, doi <https://doi.org/10.1016/j.cegh.2023.101390>*
13. *Jacob et al; "Tackling the Rising Tide: Understanding the Prevalence of Childhood Obesity in India" 28.2 (2024) Pp101-103, doi https://doi.org/10.4103/ijem.ijem_47_24*
14. *Times of India et al; "What's Behind the Rising Rates of Childhood Obesity in India?" 1.1 (2023) Pp1-2, Available from: <https://timesofindia.indiatimes.com/life-style/health-fitness/health-news/whats-behind-the-rising-rates-of-childhood-obesity-in-india/photostory/105724792.cms>*