
Dr. Amit Sachdeva & Nasim Ahmed

¹Assistant Professor, Department of Community Medicine, Indira Gandhi Medical College, Shimla, Himachal Pradesh, India. Email: dramitsachdeva2410@gmail.com

²Independent Research Scholar, Iarcon international LLP, Guwahati, Assam India. Email: nasim@iarcon.org

Received: Jan. 03, 2025; Revised: Feb. 09, 2025; Accepted: Mar. 16, 2025 ; Published: June. 28, 2025

Under a Creative Commons license Doi: 10.47310/ml.2025.v02i01.038

The Public Health Burden of Untreated Dental Caries: Global and National Perspectives

Abstract: Untreated dental caries remains a significant public health burden worldwide, affecting billions of people and contributing to pain, infections, and tooth loss. This global issue is prevalent across all age groups, with low- and middle-income countries being particularly impacted due to limited access to dental care. The consequences of untreated caries extend to diminished quality of life, school absenteeism, and economic strain. Public health strategies, including fluoride use, oral hygiene promotion, increased access to care, and addressing dietary risk factors, are essential to mitigating this burden.

Keywords: Dental caries, Public health, Oral hygiene, Fluoride, Access to care.

INTRODUCTION

Dental caries, commonly known as tooth decay or cavities, is one of the most prevalent and neglected global health issues. Affecting individuals across all age groups, untreated dental caries not only causes pain and discomfort but also leads to more serious oral health complications, including tooth loss, infections, and systemic conditions. Despite being preventable and treatable, untreated dental caries remains a significant public health burden, especially in low- and middle-income countries where access to dental care is limited.[1-4]

Worldwide, billions of people suffer from the consequences of untreated dental caries, making it a leading cause of oral health morbidity. The socioeconomic impact of dental caries is profound, contributing to school absenteeism, reduced work productivity, and financial strain due to the high costs of dental care. This article examines the global and national perspectives on the public health burden of untreated dental caries, discusses current trends and challenges, and highlights potential public health strategies aimed at reducing its impact.

Understanding Dental Caries [4-7]

1. Definition and Development of Dental Caries

Dental caries is a progressive disease caused by the breakdown of tooth enamel due to the acidic by-products of bacterial metabolism. The condition begins when bacteria in the mouth, particularly *Streptococcus mutans* and *Lactobacilli*, metabolize sugars and carbohydrates from food, producing acids that demineralize the tooth enamel. Over time, this demineralization leads to the formation of cavities.

Dental caries can affect all tooth surfaces, including the enamel, dentin, and pulp. If left untreated, the decay can spread deeper into the tooth, leading to pulpitis (inflammation of the tooth pulp), abscess formation, and, ultimately, tooth loss. Untreated caries is associated with several oral and systemic health complications, emphasizing the need for timely treatment.

2. Risk Factors for Dental Caries

How to cite this content: Amit Sachdeva; “The Public Health Burden of Untreated Dental Caries: Global and National Perspectives” Medletter, V2 I1 (2025) pp 233-237

The Public Health Burden of Untreated Dental Caries: Global and National Perspectives

Dental caries is a multifactorial disease influenced by various biological, behavioral, and environmental factors:

- **Diet:** Frequent consumption of sugary foods and beverages significantly increases the risk of dental caries. Sugars provide a substrate for bacteria, leading to acid production and demineralization of enamel.
- **Oral Hygiene:** Poor oral hygiene practices, such as inadequate brushing and flossing, allow the accumulation of plaque, a sticky biofilm of bacteria that contributes to tooth decay.
- **Fluoride Exposure:** Fluoride plays a crucial role in preventing dental caries by strengthening enamel and promoting remineralization. Inadequate fluoride exposure, whether through drinking water, toothpaste, or professional treatments, increases the risk of caries.
- **Socioeconomic Status:** Individuals from lower socioeconomic backgrounds are at higher risk for untreated dental caries due to limited access to dental care, lower health literacy, and poor dietary habits.
- **Access to Dental Care:** Limited access to preventive and restorative dental services is a significant contributor to the global burden of untreated dental caries, particularly in rural and low-income areas.

Global Burden of Untreated Dental Caries [7-10]

1. Prevalence and Incidence of Dental Caries

Dental caries is the most common non-communicable disease (NCD) globally, affecting approximately 3.5 billion people, according to recent estimates. The prevalence of untreated caries in permanent teeth is highest in adults, with nearly 2.5 billion people affected worldwide. The burden is equally substantial in children, with untreated caries in deciduous (primary) teeth affecting more than 530 million children globally.

- **Regional Variations:** The burden of dental caries varies by region, with low- and middle-income countries bearing the brunt of the disease. In sub-Saharan Africa and Southeast Asia, untreated dental caries is widespread, with limited access to preventive care and dental services exacerbating the problem. In contrast, high-income countries tend to have better access to dental care, but untreated caries remains a significant issue among underserved populations.
- **Age Distribution:** Dental caries affects individuals at every stage of life, from early childhood to old age. In children, untreated caries in deciduous teeth can lead to pain, difficulty eating, and poor nutrition, while in adults, untreated caries in permanent teeth can result in tooth loss and diminished quality of life.

2. Impact of Untreated Dental Caries on Quality of Life

The consequences of untreated dental caries extend beyond oral health, affecting general well-being and quality of life. Individuals with untreated caries often experience:

- **Pain and Discomfort:** Toothache is one of the most common symptoms of untreated dental caries, leading to difficulty eating, sleeping, and concentrating. Persistent pain can significantly impair an individual's daily functioning, particularly in children, who may miss school due to dental pain.
- **Nutritional Deficiencies:** Dental caries can affect chewing ability, leading to poor dietary choices and malnutrition. Children with untreated caries may avoid certain foods due to pain, resulting in inadequate intake of essential nutrients needed for growth and development.
- **Speech and Self-Esteem Issues:** Severe dental caries can lead to tooth loss, which can affect speech and self-confidence. Individuals, particularly adolescents and adults, may experience embarrassment due to visible decay or missing teeth, leading to social isolation and psychological distress.
- **School and Work Absenteeism:** In children, dental pain and infection are significant causes of school absenteeism, affecting their educational attainment. Similarly, adults with untreated caries may miss work due to pain or dental appointments, leading to reduced productivity and financial strain.

3. Economic Burden of Untreated Dental Caries

The economic impact of untreated dental caries is substantial, both for individuals and healthcare systems. The direct costs of treating dental caries, including fillings, crowns, root canals, and extractions, are high. In countries where dental care is not covered by national health insurance, individuals may face significant out-of-pocket expenses.

The Public Health Burden of Untreated Dental Caries: Global and National Perspectives

- **Direct Costs:** The global cost of dental treatment, including caries-related services, is estimated to exceed \$356 billion annually. These costs are disproportionately higher in high-income countries due to the increased use of advanced dental technologies and restorative treatments.

Indirect Costs: Indirect costs include lost productivity due to absenteeism from work or school, as well as the long-term consequences of untreated caries, such as tooth loss and the need for dentures or implants. In low- and middle-income countries, the indirect costs of untreated dental caries are often higher than the direct costs, as individuals are less likely to seek early treatment.

National Perspectives on the Burden of Untreated Dental Caries [1,3,11]

1. High-Income Countries

In high-income countries, such as those in North America and Europe, dental care systems are relatively well-developed, with greater access to preventive services, fluoride exposure, and education on oral hygiene. Despite this, untreated dental caries remains a significant public health issue, particularly in underserved populations, including low-income individuals, racial and ethnic minorities, and rural residents.

- **Disparities in Care:** Even in countries with advanced healthcare systems, socioeconomic disparities persist in access to dental care. In the United States, for example, dental caries is more prevalent among children from low-income families, as they are less likely to receive preventive dental services, such as sealants and fluoride treatments. Additionally, rural residents face challenges in accessing dental care due to geographic barriers and shortages of dental professionals.
- **Aging Populations:** In many high-income countries, the aging population is a growing concern for oral health. Older adults are more likely to experience untreated dental caries due to dry mouth (xerostomia) caused by medications, reduced dexterity for oral hygiene, and limited access to affordable dental care in long-term care facilities.

2. Low- and Middle-Income Countries (LMICs)

The burden of untreated dental caries is particularly high in low- and middle-income countries, where access to dental care is limited, and public health infrastructure is underdeveloped. In many LMICs, dental services are primarily available in urban centers, leaving rural populations with little to no access to preventive or restorative care.

- **Cultural and Economic Barriers:** In some regions, cultural beliefs and lack of awareness about oral health contribute to the neglect of dental care. Economic barriers, including the high cost of dental services and the absence of dental insurance, further prevent individuals from seeking care.
- **Rising Sugar Consumption:** The increasing consumption of sugary foods and beverages in LMICs, driven by globalization and urbanization, is contributing to the rising prevalence of dental caries. In countries where traditional diets were once low in sugar, the introduction of processed foods has led to a significant increase in caries rates, particularly among children.

Limited Resources for Treatment: Many LMICs lack the resources and infrastructure to provide comprehensive dental care to their populations. As a result, untreated dental caries often progresses to advanced stages, leading to pain, infection, and tooth loss. In some cases, individuals resort to traditional remedies or extractions as the only available treatment.

Public Health Strategies for Addressing the Burden of Untreated Dental Caries [12-15]

Reducing the burden of untreated dental caries requires a comprehensive approach that includes prevention, early detection, and access to affordable treatment. Public health strategies must address the root causes of dental caries, such as poor diet and inadequate oral hygiene, while also ensuring that individuals have access to the necessary dental care services.

1. Prevention Through Fluoridation

Fluoride is one of the most effective tools for preventing dental caries. Fluoridation of public water supplies has been shown to significantly reduce the incidence of caries in both children and adults. In countries with widespread water fluoridation, the prevalence of dental caries is lower than in countries without such programs.

- **Water Fluoridation:** Water fluoridation is a cost-effective public health intervention that benefits entire communities, regardless of age or socioeconomic status. It works by providing a low level of fluoride in the drinking water, which helps remineralize tooth enamel and prevent

The Public Health Burden of Untreated Dental Caries: Global and National Perspectives

decay. Despite its proven benefits, water fluoridation is not universally adopted, and some countries have faced opposition due to concerns about safety and individual choice.

- **Topical Fluoride:** In areas where water fluoridation is not feasible, alternative fluoride delivery methods, such as fluoride toothpaste, mouth rinses, and professionally applied fluoride varnishes, can help reduce the risk of caries. Public health campaigns promoting the use of fluoride toothpaste, especially in children, are essential for improving oral health outcomes.

2. Promoting Oral Hygiene Practices

Good oral hygiene practices, such as regular brushing and flossing, are critical for preventing dental caries. Public health programs should focus on educating individuals about the importance of oral hygiene and providing access to affordable oral care products.

- **School-Based Programs:** School-based oral health programs that teach children how to brush and floss properly can have a lasting impact on their oral health. In addition to education, these programs often provide free toothbrushes, toothpaste, and fluoride treatments to children in underserved areas. Such programs are particularly effective in reducing the prevalence of caries in low-income populations.
- **Community Outreach:** In rural and underserved communities, mobile dental clinics and community health workers can provide basic oral health services and education. These programs can help reach populations that may not have regular access to dental care and raise awareness about the importance of preventive measures.

3. Expanding Access to Dental Care

Ensuring that individuals have access to affordable dental care is essential for reducing the burden of untreated dental caries. Public health policies should focus on expanding coverage for dental services, particularly for vulnerable populations, such as children, low-income individuals, and the elderly.

- **Integrating Oral Health into Primary Care:** Integrating oral health services into primary healthcare settings can help improve access to dental care. In many countries, oral health is treated separately from general health, which limits opportunities for early detection and treatment. By training primary care providers to conduct basic oral health screenings and refer patients to dental specialists, more individuals can receive timely care.
- **Public-Private Partnerships:** Partnerships between governments, non-governmental organizations (NGOs), and the private sector can help improve access to dental care in underserved areas. These partnerships can provide funding for mobile dental clinics, free or low-cost dental services, and public health campaigns aimed at preventing dental caries.

4. Addressing Dietary Risk Factors

Reducing sugar consumption is a key strategy for preventing dental caries. Public health campaigns that promote healthy eating habits and limit the intake of sugary foods and beverages can help reduce the incidence of caries, particularly in children.

- **Sugar Taxation:** Several countries have implemented taxes on sugary beverages as a way to reduce consumption and generate revenue for public health programs. Research has shown that sugar taxes can lead to a significant decrease in the consumption of sugary drinks, which are a major contributor to dental caries.
- **Nutrition Education:** Public health programs should include nutrition education that emphasizes the importance of a balanced diet low in sugar. Schools, community centers, and healthcare facilities can provide resources and guidance on healthy eating to help individuals make informed dietary choices.

CONCLUSION

The public health burden of untreated dental caries is a significant global issue, affecting billions of people and contributing to pain, discomfort, and diminished quality of life. While the burden is felt most acutely in low- and middle-income countries, high-income countries also face challenges in ensuring equitable access to dental care.

Preventing and treating dental caries requires a multifaceted approach that includes promoting fluoride use, improving oral hygiene practices, expanding access to affordable dental care, and addressing the dietary factors that contribute to caries. Public health strategies must prioritize vulnerable populations, such as children, low-income individuals, and rural residents, to reduce disparities in oral health outcomes.

The Public Health Burden of Untreated Dental Caries: Global and National Perspectives

By investing in preventive measures and ensuring that individuals have access to timely and affordable treatment, we can significantly reduce the global burden of untreated dental caries and improve the overall health and well-being of populations worldwide.

REFERENCES

1. Nath et al; "The Global Prevalence and Severity of Dental Caries among Racially Minoritized Children: A Systematic Review and Meta-Analysis" *Caries Research* 57.4 (2023) Pp 485-508, doi <https://doi.org/10.1159/000534371>
2. World Health Organization; "WHO Highlights Oral Health Neglect Affecting Nearly Half of the World's Population" WHO (2022) Pp n/a, cited 2024 Oct 14, available from <https://www.who.int/news/item/18-11-2022-who-highlights-oral-health-neglect-affecting-nearly-half-of-the-world-s-population>
3. Kassebaum et al; "Global Burden of Untreated Caries: A Systematic Review and Metaregression" *Journal of Dental Research* 94.5 (2015) Pp 650-658, doi <https://doi.org/10.1177/0022034515573272>
4. Qin et al; "Changes in the Global Burden of Untreated Dental Caries From 1990 to 2019: A Systematic Analysis for the Global Burden of Disease Study" *Heliyon* 8.9 (2022) Pp e10714, doi <https://doi.org/10.1016/j.heliyon.2022.e10714>
5. Aditya Birla Memorial Hospital; "Demystifying Dental Caries: Understanding Causes, Prevention, and Treatment" Aditya Birla Memorial Hospital (2023) Pp n/a, cited 2024 Oct 14, available from <https://adityabirlahospital.com/blogs/Demystifying-Dental-Caries-Understanding-Causes-Prevention-and-Treatment>
6. Omega Hospitals; "Understanding Dental Caries: Causes, Symptoms, Prevention, and Treatment" Omega Hospitals (2023) Pp n/a, cited 2024 Oct 14, available from <https://www.omegahospitals.com/blog/understanding-dental-caries-causes-symptoms-prevention-and-treatment/>
7. Cleveland Clinic; "Cavities: Symptoms, Causes, Diagnosis, Treatment" Cleveland Clinic (2023) Pp n/a, cited 2024 Oct 14, available from <https://my.clevelandclinic.org/health/diseases/10946-cavities>
8. Mayo Clinic; "Cavities: Symptoms and Causes" Mayo Clinic (2023) Pp n/a, cited 2024 Oct 14, available from <https://www.mayoclinic.org/diseases-conditions/cavities/symptoms-causes/syc-20352892>
9. Fejerskov; "Concepts of Dental Caries and Their Consequences for Understanding the Disease" *Community Dentistry and Oral Epidemiology* 25.1 (1997) Pp 5-12, doi <https://doi.org/10.1111/j.1600-0528.1997.tb00894.x>
10. Giacaman et al; "Understanding Dental Caries as a Non-Communicable and Behavioral Disease: Management Implications" *Frontiers in Oral Health* 3 (2022) Pp 764479, doi <https://doi.org/10.3389/froh.2022.764479>
11. Bernabe et al; "Global, Regional, and National Levels and Trends in Burden of Oral Conditions From 1990 to 2017: A Systematic Analysis for the Global Burden of Disease 2017 Study" *Journal of Dental Research* 99.4 (2020) Pp 362-373, doi <https://doi.org/10.1177/0022034520908533>
12. Chen et al; "Oral Health Policies to Tackle the Burden of Early Childhood Caries: A Review of 14 Countries/Regions" *Frontiers in Oral Health* 2 (2021) Pp 670154, doi <https://doi.org/10.3389/froh.2021.670154>
13. Lee; "Diagnosis and Prevention Strategies for Dental Caries" *Journal of Lifestyle Medicine* 3.2 (2013) Pp 107-109
14. Duangthip et al; "Challenges in Oral Hygiene and Oral Health Policy" *Frontiers in Oral Health* 1 (2020) Pp 575428, doi <https://doi.org/10.3389/froh.2020.575428>
15. Bogale et al; "Dental Caries Experience and Associated Factors in Adults: A Cross-Sectional Community Survey Within Ethiopia" *BMC Public Health* 21 (2021) Pp 180, doi <https://doi.org/10.1186/s12889-021-10199-6>